Parent	
Baby	
DOB	_ Time
Birthweight	

BREASTFEEDING RECORD



	DATE	6	7	8	9	10	11	noon	1	2	3	4	5	6	7	8	9	10	11	mid	1	2	3	4	5	#	#	#
		am	am	am	am	am	am		pm	pm	pm	pm	pm	pm	pm	pm	pm	pm	pm	night	am	am	am	am	am	feed	U	S
1																												
	Generally	: feed	on de	mand	SO	me ba	bies s	leep lon	ig peri	ods o	n the f	irst da	ay – ta	ke ad	vantag	ge by g	getting	g ahea	ad on	your ow	n slee	р	11					
2																												
	Generally	: baby	/ likely	to be	fussy	while	waitin	g for mil	lk to c	ome ii	n ke	ep fee	eding	on der	mand	at leas	st 8x p	er da	y u	se nippl	e crea	m r	no soot	thers of	or form	ula		
3																												
	Generally	: feed	on de	mand	, wake	e a sle	epy ba	aby to fe	ed mi	nimal	ly evei	ry 3ho	urs; if	jaund	iced, s	sunba	the na	ked ir	n wind	ow; pos	sible u	irate c	rystals	in dia	per			
4																												
	Generally	: deal	with v	ery ful	ll brea	sts	if eng	orged, p	oump	or har	nd exp	ress to	o softe	en are	ola be	fore fe	eeds,	use co	old cal	bbage le	aves	after fe	eeds					
5																												
	Generally	: watc	h for c	hange	e in sto	ool col	or to y	ellow, a	and inc	rease	e in an	nount	of urir	ie							-							
6																												
	Generally	/: baby	/ much	ı less i	fussy	once g	getting	creamy	/ brea	stmilk																•		
7																												
	Generally	/: gaini	ing 30	g (= 1	ounce	e) or m	nore pe	er day																				

BF	Breastfeed	PLAN:
EBM	Expressed Breast Milk	
F	Formula	
U	Urine	
S	Stool	



Steps to ensure a good milk supply & healthy baby in the first weeks

1. Feed on demand

- 2. Unwrap, waken & feed a sleepy baby every 2 1/2 to 3 hours
- 3. Pay attention to latch
 - a. Latch baby with a big, open mouth like a yawn
 - b. Relatch whenever a bad latch is noticed
 - c. Any discomfort should not last more than a minute
- 4. Encourage baby to be awake & vigorous at the breast
 - a. Feed skin to skin
 - b. Stimulate a sleepy baby tickling, blowing, cold cloth
 - c. Use breast compressions if necessary [see video at <u>www.drjacknewman.com</u>]
- 5. Exclusively breastfeed (NO water, tea, juice, formula, soothers or fingers)
- 6. Take good care of mom eat and drink well, get as much rest as possible

Normal newborn

٨	Temperatu	re 36.5-37.5°C
	0	Warm body, cool hands & feet
	0	Pink blue/purple hands & feet in first day
≻	Breathing:	
	0	Irregular breathing pattern
	0	30-60 respirations per minute
	0	Apnea up to 20 seconds
۶	Skin:	
	0	Jaundice/yellow after first 24 hours, not extending downwards past chest or into eyes
	0	Newborn acne on face/body &/or milia (white spots on face)
۶	Input:	
	0	Feeding minimum 8x/day, vigorous sucking at least 20 mins
	0	Occasional cluster feeding
۶	Output:	
	0	Urine: 1 for every day of life until milk comes in 4+ heavy diapers after milk comes in
	0	Urate crystals (pink/orange color) in urine before milk comes in
	0	Stool: Meconium in first few days transitional green breastmilk yellow, liquid or pasty
۶	Weight:	
	0	Weight loss up to 10% of birthweight
	0	Regain of birthweight by 2-3 weeks
	0	Average of 1 oz (30 g) per day once gaining